

MONITORING HYDROLOGICAL REGIME OF THE EUPHRATES-TIGRIS RIVER BASIN USING SATELLITE OBSERVATIONS

Elena A. Zakharova,
Alexei V. Kouraev,
Igor Polikarpov,
Faiza Al-Yamani,
Jean-François Crétaux

Laboratoire d'Etudes en Geophysique et Oceanographie Spatiales (LEGOS), Toulouse France
State Oceanography Institute, St. Petersburg branch, St. Petersburg, Russia
Mariculture and Fishery Dept., Kuwait Institute for Scientific Research, Kuwait



CENTRE NATIONAL D'ETUDES SPATIALES


E-T WATERSHED

A = 765 000 km²

W=105 km³

L_{Euphrates} = 2800 km
W_{Euphrates} = 30 km³/y

L_{Tigris} = 1750 km
W_{Tigris} = 55 km³/y



E-T WATERSHED

4 riparian states:

Turkey, Syria, Iran, Iraq

Before 70s

Turkey by Euphrates

31 km³/y

Syria

27 km³/y

Iraq

? km³/y

Persian Gulf

From 90s

Gap project in Turkey
+ 22 reservoirs
+ 1.7 mln ha in irrigation

16 km³/y


Syria

9 km³/y

Iraq

? km³/y

Influence on fishery
resources in the
Persian Gulf



E-T WATERSHED

Main reservoirs

Tigris R.

Mosul W = 13.5 km³

Euphrates R.

Keban W = 31 km³

Karakaya W = 10 km³

Ataturk W = 49 km³


Tharthar Lake

W = 73 km³

40 reservoirs

Water storage = 5 times
of annual ET discharge

In situ data -
scarce and
difficult to access



DATA AQUISITION


T/P	(Oct 1992- Aug 2002, new orbit - Aug 2002-2005)
Jason-1	(since Feb 2002)
GFO	(since Jan 2000)
ENVISAT	(since Nov 2002)
GRACE	(data since 2003)

FOOTPRINT: RADAR - 12 km (ocean), RADIOMETER - 22 TO 42 km, GRACE ~ 625 km²

REPEAT PERIOD: 10 days (T/P, Jason), 17 days (GFO) and 35 (ENVISAT),

GRACE product: maps of 1-degree-square mean geoid height differences between 30-day solutions and EIGEN_GL04S static solution (10 days resolution).


From International Gravimetric Bureau (<http://bgi.cnes.fr:8110/>)



ENVISAT ground tracks and selected altimetric measures for all cycles (yellow circles) over the upper part of the Ataturk water reservoir. Background - Landsat TM image from ca 1990.

RESERVOIRS: WATER LEVEL


Temporal changes of reservoir level (mm) from altimetric data



Altimetric level - relative to geoid

T/P + GFO + ENVISAT: Karakaya, Mosul and Tharthar




Cross-comparison of satellites Times series for 11 reservoirs



RESERVOIRS: WATER VOLUME

Preliminary assessment (surface=const)

1999 - 2002
30 - 40 % of volume loss in
reservoirs on Euphrates R.





MESOPOTAMIAN MARSHES

Changes in the marshes from Landsat
MSS imagery

20,000 km² in 70s

3,000 km² in 90s




Multi-Spectral Scanner (MSS) aboard Landsat mom <http://landsat.gsfc.nasa.gov/>




MARSHES


SPATIAL AND TEMPORAL VARIABILITY




Landsat image ca 2000



ENVISAT Tr.197



ENVISAT Tr.098



- -1000 to 0
- 0 to 1000
- 1000 to 1500
- 1500 to 2000
- 2000 to 3000
- 3000 to 4200
- 4200 to 6000


Backscatter in Ku band,
ice1 algorithm,
100*dB

EASTERN MARSHES



- T/P 31



Landsat image ca 1990




Landsat image ca 2000



- A - southern marshes
- B - confluence of Tigris and Euphrates
- C - part of eastern marshes, drained out between 3 and 23 June 1996
- D - eastern marshes




MARSHES - INONDATION

ENVISAT Tr. 642



Landsat image ca 2000


Backscatter in Ku band,
ice1 algorithm,
100*dB




- A - river channel
- B - flooded reservoir
- C - large flooded area
- D - partly flooded central marshes
- E - southern marshes have been drained out in 1996 and were flooded in 2004
- F - channel

MARSHES - INONDATION

ENVISAT 741




Landsat image ca 2000




A B C D E F G

Backscatter in Ku band,
ice1 algorithm,
100*dB



MARSHES - INONDATION

along track seasonal variability




*Landsat image ca 2000
and Envisat tracks*

$$\% \text{ of submerged lands} = \frac{\text{number of points with } \sigma_0 \text{ Ku} \geq 20 \text{ db}}{\text{total number of points}}$$

Perspectives:


- calibration of flooding areas with Modis
- estimation of water storage using altimetry





GRACE MISSION:

VARIABILITY IN WATER RESOURCES


Watershed surface
765 000 km²



Gravity anomalies (mm) over the
TE basin
for 2004





RELATION BETWEEN OBSERVED PARAMETERS



Cross-correlation coefficients between the part of submerged lands and other parameters , n≈40

Time lag, days	0	10	20	30	40	50	60	70	80	90	100	110	120
GRACE, ET	0.58	0.63	0.72	0.78	0.78	0.80	0.79	0.64	0.65	0.60	0.49	0.29	0.24
GRACE, Iran	0.68	0.75	0.78	0.83	0.80	0.73	0.69	0.67	0.61	0.52	0.42	0.25	0.12
GRACE, Turkey	0.56	0.57	0.65	0.73	0.73	0.73	0.71	0.58	0.59	0.54	0.38	0.18	0.12
GRACE, Low Iraq	0.56	0.66	0.69	0.75	0.78	0.73	0.77	0.69	0.63	0.57	0.54	0.40	0.32
Altimetry, dV in all reservoirs	0.20	0.29	0.38	0.47	0.55	0.63	0.69	0.74	0.77	0.79	0.79	0.77	0.72

IMPACT ON PERSIAN GULF



FUTURE WORK

- * **Assessment of reservoirs surface-level relationship using optical imagery**
- * **More precise estimation of reservoirs volumetric changes**
- * **Altimetry + optical imagery - flooded regions**
- * **Surface and level of flooded zones**
- * **Estimation of annual and seasonal changes in water storage**
- * **Freshwater input to the Persian gulf**