

ERS Precision Orbit Determination and Accuracy Verification

C. Shum, J. Ries, J. Bordi, J. Seago, and B. Tapley

Center for Space Research, The University of Texas at Austin,
Austin, Texas 78712, USA. Tel: 512-471-5573, Internet:
shum@csr.utexas.edu

Development in the Earth's gravity field and ocean tide models, non-conservative force models, terrestrial reference frame definition, and precision orbit determination methodologies to employ dual-satellite altimeter crossovers, has enable the current radial orbit accuracy for ERS-1 and ERS-2 to approach the 5 cm rms level. These improved orbit determination methodologies enable an improvement in both the constant and the variable component of the ERS-1 and ERS-2 orbit error, and provide improved alignment of the absolute sea surface measurements from ERS within the terrestrial reference frame. Non-conservative forces are the current limitations for the ERS-1 orbit accuracy. The significant contribution by the global PRARE range and range-rate tracking measurements provides an improvement in the handling of non-conservative forces for ERS-2. In addition to crossover analyses, independently determined orbits were used to assess and verify the radial accuracy of the orbits. The orbit accuracy assessment also employed a technique which uses TOPEX/POSEIDON measurements for the characterization of the geographically correlated component of the gravity error in ERS-1 and ERS-2 orbits.