

ReadMe-first Technical Note

(RM-TN)

Document Version

| Version | Date | Description | Author |
|---------|------------|--|--|
| 1.6 | 13/11/2023 | Refers to CryoSat-2 SMOS Merged Product version v206 | Stefan Hendricks (AWI) Lars Kaleschke (AWI) |
| 1.5 | 08/11/2022 | Refers to CryoSat-2 SMOS Merged Product version v205 | Stefan Hendricks (AWI) Lars Kaleschke (AWI) |
| 1.4 | 01/11/2021 | Refers to CryoSat-2 SMOS Merged Product version v204 | Stefan Hendricks (AWI) |
| 1.3 | 07/10/2020 | Refers to CryoSat-2 SMOS Merged Product version v203 | Robert Ricker (AWI) |
| 1.2 | 27/09/2019 | Refers to CryoSat-2 SMOS Merged Product version v202 | Robert Ricker (AWI) |
| 1.1 | 19/10/2018 | Revision after comments from ESA | Robert Ricker (AWI) |
| 1.0 | 15/06/2018 | Draft of the ReadMe-first Technical Note | Robert Ricker (AWI) |

Applicable Documents

| Abbreviation | Name | Description |
|--------------|---------------------------|--------------------------------------|
| ATBD | AWI_ESA_CS2SMOS_ATBD_v2.5 | Algorithm Theoretical Basis Document |
| PDD | AWI_ESA_CS2SMOS_PDD_v1.6 | Product Description Document |

| | |
|--|---|
| Read-me-first note for the release of theCryoSat-2/SMOS Merged Sea Ice Thickness (CS2SMOS) | |
| Product version | v206 |
| Release date by ESA | 13.11.2023 |
| Author(s) | Robert Ricker, Stefan Hendricks, Lars Kaleschke |
| Further information | <p>A detailed description of the processing algorithm can be found in the Algorithm Theoretical Basis Document (ATBD). Information about the data product ca be found in the Product description document (PDD):</p> <p>The documents are available here: https://earth.esa.int/eogateway/catalog/smos-cryosat-l4-sea-ice-thickness</p> <p>Information on how to access the CS2SMOS ice thickness data from AWI can be found here: https://spaces.awi.de/confluence/x/DwVmEQ</p> <p>Information on how to access the CS2SMOS ice thickness data from ESA can be found here: https://smos-diss.eo.esa.int/oads/access/</p> |
| How to cite the data | <p>1. Please cite:</p> <p>Data set ESA/AWI CryoSat/SMOS L4 sea ice thickness, https://doi.org/10.57780/sm1-4f787c3</p> <p>Ricker, R., Hendricks, S., Kaleschke, L., Tian-Kunze, X., King, J., and Haas, C.: A weekly Arctic sea-ice thickness data record from merged CryoSat-2 and SMOS satellite data, The Cryosphere, 11, 1607-1623, https://doi.org/10.5194/tc-11-1607-2017, 2017.</p> <p>2. Include the following phrase into the acknowledgment:</p> <p>“The production of the merged CryoSat-SMOS sea ice thickness data was funded by the ESA project SMOS & CryoSat-2 Sea Ice Data Product Processing and Dissemination Service, and data of version 2.06 from DATE to DATE were obtained from (AWI or ESA).”</p> |
| Contact for helpline | For all issues related to data access, please contact ESA’s HelpDesk at eohelp@esa.int |

| | |
|---|---|
| Comments to CryoSat-2/SMOS merged product | For questions and feedback, please contact: cs2smos-support@awi.de |
|---|---|

1. Introduction

The read-me-first note provides information about improvements with regard to the previous releases, data caveats, and instruction about how to use auxiliary data and uncertainties contained in the product.

The current product version v206 is available from October 2023 to present. It will replace v205, after reprocessing is completed. The product is continuously generated within the framework of the ESA project **SMOS & CryoSat-2 Sea Ice Data Product Processing and Dissemination Service**. The product is only available for the Northern Hemisphere from October to April.

2. Main Improvements and Changes in the current Data Set

The main changes in v206 compared to previous version v205 are algorithm updates of the CryoSat-2 source sea ice thickness data sets:

CryoSat-2 version 2.6 update

- Reprocessing with CryoSat-2 ICE Level-1b algorithm baseline-E (in v2.5, baseline-E data was only used since Oct 2020).
- Use the OSI-SAF / C3S sea ice type (interim) climate data record. Notable changes in sea ice type information in the first half of October.
- There had been cases of incomplete CryoSat-2 near real-time L1B data on the production system in v2.5. The latency of near real-time production has been increased from 36h to 48h

CryoSat-2/SMOS data fusion update

None

3. Product Performance

CS2SMOS, CryoSat-2 and SMOS Ice Thickness Uncertainties

The uncertainties of the CryoSat-2 and SMOS sea ice thickness observations are crucial for the data merging and the interpolation. Figure 1 shows the relative uncertainties of CryoSat-2 and SMOS for November 2013 and April 2014. While the SMOS relative uncertainties are lowest for very thin ice, CS2 relative thickness uncertainties are smaller over thick ice and rise asymptotic towards thickness values < 1 m, which is due to the different methodical approach.

The merged product (CS2SMOS) takes advantage of the complementary uncertainties. CS2SMOS merged ice thickness shows a significant reduction in the relative uncertainty with regard to the thickness uncertainties of the thin ice in the CryoSat-2 product, and the thick ice in the SMOS product.




Figure 1: Binned Relative uncertainties for November 2013 and April 2014. Error bars indicate the standard deviation of relative uncertainties within the 10 cm bin width.

Product validation

For validation, we use ULS ice draft data from moorings that have been deployed at four different sites in the Beaufort Sea within the BGEF project (Woods Hole Oceanographic Institution (WHOI), Krishfield and Proshutinsky, 2006). Data are sampled at 2s intervals. Ice draft is converted to ice thickness by multiplying the draft by 1.1 (Rothrock et al., 2008). Table 1 provides the position of the moorings (A, B, D) in the Beaufort Sea and information about the ULS record periods. For the comparison of the merged CS2SMOS ice thickness and ULS ice thickness measurements, the data was averaged over 24 hours to obtain the daily mean effective ice thickness for each ULS. In the last step, daily retrievals are averaged weekly on

a 25 km EASE2 grid to cover the same period as the weekly CS2SMOS products. Since the positions of the moorings are steady, one data point for each ULS is retrieved per week and is then compared with the weekly mean of the CS2SMOS ice thickness. Figure 2 shows the positions of the moorings and the corresponding differences to the CS2SMOS ice thickness. Mean differences (MD) are calculated by subtracting satellite ice thickness from the ULS ice thickness. Considering the entire ULS data set as the reference, the mean difference is 3.5 cm, while the root mean square deviation (RMSD) is 0.29 m. This shows an improvement compared to the previous product versions (Table 2).

Table 1: Mooring sites with ULS measurements used in this document.

| Mooring Site | ULS record periods | Location |
|--------------|---|----------------|
| A | 08/2003 – 10/2016 | 150.0°W 75.0°N |
| B | 08/2003 – 09/2005 09/2006 – 09/2009 10/2010 – 10/2016 | 150.0°W 80.0°N |
| D | 09/2006 – 10/2016 | 140.0°W 74.0°N |

Table 2: Product validation with BGEP ULS measurements for different CS2SMOS product versions: mean differences (MD) and root mean square deviation (RMSD). Please note that the comparison periods changed over time. The first official ESA release was version v201.

| Product version (release date) | MD [m] | RMSD [m] |
|--------------------------------|--------|----------|
| v200 | 0.1 | 0.36 |
| v201 (2018) | 0.02 | 0.34 |
| v202 (2019) | -0.079 | 0.322 |
| v203 (2020) | -0.067 | 0.316 |
| v204 (2021) | -0.078 | 0.316 |
| v205 (2022) | 0.05 | 0.302 |
| v206 (2023) | 0.035 | 0.291 |




Figure 2: Example CS2SMOS sea ice thickness map (22-28 Oct. 2022) and ULS locations (left panel), and scatter plot between ULS and CS2SMOS v205 ice thickness over the period Nov 2010 – Oct 2022. The RMSD represents the root mean square deviation. MD represents the mean difference between CS2SMOS v206 and ULS ice thickness.

4. Known Issues

Major caveats are listed below:

| Issue | Product Version | Status |
|--|-----------------|--|
| Underestimation of SMOS ice thickness when ice concentration is lower than 100% | 206 | Open |
| Fundamental calibration of CryoSat- 2 range retracking algorithm required | 206 | Open |
| Sea ice classification (first-year / multi-year ice) auxiliary data not available in first half of October in each year with an impact on CryoSat-2 sea ice thickness. This results in a biased background field for reprocessed CS2SMOS data that uses the C3S sea ice type (interim) climate data record v2.0 for each October in the data record. This issue is no longer present in the next version of the C3S sea ice type (interim) climate data record (v3.0) that is scheduled for release in early 2023. | 206 | Closed (included in CryoSat-2 v2.6) |

5. Future algorithm evolution

We want to investigate the capability of expanding the processing to the Southern hemisphere. Moreover, we will aim to improve the retrieval algorithm. In particular, we will investigate the possibility to apply a multiyear ice concentration product to better distinguish between first-year and multiyear sea ice.

References

- Kaleschke, L., Tian-Kunze, X., Maaß, N., Beitsch, A., Wernecke, A., Miernecki, M., Müller, G., Fock, B. H., Gierisch, A. M., Schlünzen, K. H., Pohlmann, T., Dobrynin, M., Hendricks, S., Asseng, J., Gerdes, R., Jochmann, P., Reimer, N., Holfort, J., Melsheimer, C., Heygster, G., Spreen, G., Gerland, S., King, J., Skou, N., Søbjaerg, S. S., Haas, C., Richter, F., and Casal, T.: SMOS sea ice product: Operational application and validation in the Barents Sea marginal ice zone, *Remote Sensing of Environment*, 180, 264-273, doi:<http://dx.doi.org/10.1016/j.rse.2016.03.009>, special Issue: ESA's Soil Moisture and Ocean Salinity Mission - Achievements and Applications, 2016.
- Krishfield, R., Proshutinsky, A., 2006. BGOS ULS Data Processing Procedure. Woods Hole Oceanographic Institution, March 2006, 14 PP.
- Ricker, R., Hendricks, S., Kaleschke, L., Tian-Kunze, X., King, J., and Haas, C.: A weekly Arctic sea-ice thickness data record from merged CryoSat-2 and SMOS satellite data, *The Cryosphere*, 11, 1607-1623, <https://doi.org/10.5194/tc-11-1607-2017>, 2017.
- Ricker, R., Hendricks, S., Girard-Ardhuin, F., Kaleschke, L., Lique, C., Tian-Kunze, X., Nicolaus, M., Krumpen, T. (2017). Satellite-observed drop of Arctic sea ice growth in winter 2015-2016. *Geophysical Research Letters*, 44(7), 3236-3245.
- Ricker, R., Hendricks, S., Helm, V., Skourup, H., and Davidson, M.: Sensitivity of CryoSat-2 Arctic sea-ice freeboard and thickness on radar-waveform interpretation, *The Cryosphere*, 8, 1607–1622, doi:10.5194/tc-8-1607-2014, 2014.
- Rothrock, D. A., Percival, D. B. and Wensnahan, M.: The decline in arctic sea-ice thickness: Separating the spatial, annual, and interannual variability in a quarter century of submarine data, *Journal of Geophysical Research: Oceans*, 113(C5), 2008.
- Tian-Kunze, X., Kaleschke, L., Maaß, N., Mäkynen, M., Serra, N., Drusch, M., and Krumpen, T.: SMOS-derived thin sea ice thickness: algorithm baseline, product specifications and initial verification, *The Cryosphere*, 8, 997–1018, doi:10.5194/tc-8-997-2014, URL <http://www.the-cryosphere.net/8/997/2014/>, 2014.
- Wingham, D., Francis, C., Baker, S., Bouzinac, C., Brockley, D., Cullen, R., de Chateau-Thierry, P., Laxon, S., Mallow, U., Mavrocordatos, C., Phalippou, L., Ratier, G., Rey, L., Rostan, F., Viau, P., and Wallis, D.: CryoSat: A mission to determine the fluctuations in Earth's land and marine ice fields, *Advances in Space Research*, 37, 841 – 871, doi:10.1016/j.asr.2005.07.027, 2006.